

Quantifying uncertainty due to fission-fusion dynamics as a component of social complexity

Journal:	<i>Proceedings B</i>
Manuscript ID	RSPB-2018-0532
Article Type:	Research
Date Submitted by the Author:	14-Mar-2018
Complete List of Authors:	Ramos-Fernandez, Gabriel; Instituto Politecnico Nacional, CIIDIR Unidad Oaxaca; Universidad Nacional Autonoma de Mexico, Centro de Ciencias de la Complejidad King, Andrew; Swansea University, Biosciences Beehner, Jacinta; University of Michigan, Department of Psychology; University of Michigan, Department of Anthropology Bergman, Thore; University of Michigan, Psychology; University of Michigan, Ecology and Evolutionary Biology Crofoot, Margaret; University of California, Davis, Anthropology; Smithsonian Tropical Research Institute Difiore, Anthony; University of Texas at Austin, Department of Anthropology Lehmann, Julia; University of Roehampton, Department of Life Sciences Schaffner, Colleen; Universidad Veracruzana, Instituto de Neuroetologia Snyder-Mackler, Noah; University of Washington, Department of Psychology; University of Washington, Center for Studies in Demography and Ecology ZUBERBÜHLER, Klaus; University of Saint Andrews, Department of Psychology and Neuroscience; Institute of Biology, University of Neuchatel, Department of Comparative Cognition Aureli, Filippo; Universidad Veracruzana, Instituto de Neuroetologia; Liverpool John Moores University, Research Centre in Evolutionary Anthropology and Palaeoecology Boyer, Denis; Universidad Nacional Autonoma de Mexico, Instituto de Fisica; Universidad Nacional Autonoma de Mexico, Centro de Ciencias de la Complejidad
Subject:	Behaviour < BIOLOGY, Cognition < BIOLOGY, Ecology < BIOLOGY
Keywords:	fission-fusion dynamics, social complexity, social uncertainty, social cognition, social intelligence hypothesis, Shannon's entropy
Proceedings B category:	Behaviour



Quantifying uncertainty due to fission-fusion dynamics as a component of social complexity

Gabriel Ramos-Fernandez^{1,2}, Andrew J. King³, Jacinta C. Beehner⁴, Thore J. Bergman⁵, Margaret C. Crofoot^{6,7}, Anthony Di Fiore⁸, Julia Lehmann⁹, Colleen M. Schaffner¹⁰, Noah Snyder-Mackler¹¹, Klaus Zuberbühler^{12,13}, Filippo Aureli^{10,14}, and Denis Boyer^{15,2}

¹*Instituto Politécnico Nacional, CIIDIR Unidad Oaxaca, Hornos 1003, Santa Cruz Xoxocotlán, Oaxaca, 71230, Mexico*

²*Centro de Ciencias de la Complejidad, Universidad Nacional Autónoma de México, Mexico*

³*Department of Biosciences, College of Science, Swansea University, UK*

⁴*Departments of Psychology and Anthropology, University of Michigan, USA*

⁵*Departments of Psychology and Ecology and Evolutionary Biology, University of Michigan, USA*

⁶*Department of Anthropology, University of California, USA*

⁷*Smithsonian Tropical Research Institute, Panama*

⁸*Department of Anthropology, University of Texas at Austin, USA*

⁹*Department of Life Sciences, Roehampton University, UK*

¹⁰*Instituto de Neuroetología, Universidad Veracruzana, Mexico*

¹¹*Department of Psychology and Center for Studies in Demography and Ecology, University of Washington, USA*

¹²*School of Psychology and Neuroscience, University of St Andrews, Scotland, UK*

¹³*Department of Comparative Cognition, Institute of Biology, University of Neuchâtel, Switzerland*

¹⁴*Research Centre in Evolutionary Anthropology and Palaeoecology, Liverpool John Moores University, UK*

¹⁵*Instituto de Física, Universidad Nacional Autónoma de México, Mexico*

Abstract

Groups of animals (including humans) may show flexible grouping patterns, in which temporary aggregations or subgroups come together and split, changing composition over short temporal scales, (*i.e.* fission and fusion). A high degree of fission-fusion dynamics may constrain the regulation of social relationships, introducing uncertainty in interactions between group members. Here we use Shannon's entropy to quantify the predictability of subgroup composition for three species known to differ in the way their subgroups come together and split over time: spider monkeys (*Ateles geoffroyi*), chimpanzees (*Pan troglodytes*) and geladas (*Theropithecus gelada*). We formulate a random expectation of entropy that considers subgroup size variation and sample size, against which the observed entropy in subgroup composition can be compared. Using the theory of set partitioning, we also develop

12 a method to estimate the number of subgroups that the group is likely to be divided into,
13 based on the composition and size of single focal subgroups. Our results indicate that
14 Shannon's entropy and the estimated number of subgroups present at a given time provide
15 quantitative metrics of uncertainty in the social environment (within which social relation-
16 ships must be regulated) for groups with different degrees of fission-fusion dynamics. These
17 metrics also represent an indirect quantification of the cognitive challenges posed by socially
18 dynamic environments. Overall, our novel methodological approach provides new insight
19 for understanding the evolution of social complexity and the mechanisms to cope with the
20 uncertainty that results from fission-fusion dynamics.

21 *Keywords:* fission-fusion dynamics, social complexity, social uncertainty, social cognition,
22 social intelligence, Shannon's entropy

23 **1 Introduction**

24 Fission-fusion dynamics are a property of any social system that displays temporal variation in
25 cohesion, subgroup size and composition [1]. These dynamics have been shown to be adaptive,
26 especially for species that forage on heterogenous resources, since they afford individuals the
27 opportunity to adjust subgroups to current and local resource abundance [2–5]. The fluid nature
28 of subgroup composition due to a high degree of fission-fusion dynamics generates a complex
29 environment within which social relationships must be regulated and consequently, constitutes
30 a potential selective pressure for cognitive abilities required to keep track of interactions in
31 frequently changing social settings [1, 6].

32 Given the relevance and widespread occurrence of fission-fusion dynamics across taxa, it is nec-
33 essary to have metrics which can capture the variability in fission-fusion dynamics within and
34 across species and environments [1]. A high degree of fission-fusion dynamics, where subgroup
35 composition is frequently changing, increases the diversity of contexts in which the same indi-
36 viduals interact, making it more difficult to track social information in species where individual
37 recognition exists [7]. While several studies quantifying social complexity deal with the diversity
38 of relationships that individuals hold [8–10], quantifying the diversity of contexts in which these

relationships are established and maintained can also be useful as a measure of social complexity [11]. To our knowledge there are no quantitative measures of this diversity of contexts for social interaction [12]. Here we propose such a measure based on information theory.

When fission-fusion dynamics occur within the boundaries of a larger, stable group, subgroups can be thought of as subsets of individuals taken from a finite set. Thus, information entropy or information content (hereafter Shannon's entropy; [13]) is an ideal measure for the extent to which subgroup composition is predictable, because the measure was derived precisely for a process in which discrete symbols are selected from a finite set. Suppose that a group of $n = 10$ individuals can be found divided in subgroups. The total number of different subgroups that can be formed, S , is $2^n - 1 = 1023$. If, for instance, all subgroups are equally likely, we would have a $1/1023$ chance of guessing the correct composition of a subgroup chosen at random. For such a set of possible subgroup compositions, the information content is equal to $\log_2(S) \simeq 10$ (in bits), which is the average minimal number of yes/no questions needed to figure out the composition of a randomly chosen subgroup. When all the S subgroup compositions are equally likely, the dataset's information content is maximal. More generally, knowing the probability p_i of observing a subgroup with composition i (with i an index ranging from 1 to S), the Shannon's entropy or information content H , for all possible subgroups S with associated probabilities p_i is:

$$H = - \sum_{i=1}^S p_i \log_2 p_i \quad (1)$$

If all possible subgroup compositions are observed with similar frequencies, H will be near maximal, implying that each observed composition carries a high information content. In contrast, if some subgroup compositions become more likely than others, our uncertainty will decrease (and so will H), i.e. on average, the observation of a particular subgroup composition will reveal less information.

Shannon's entropy is therefore directly related to the uncertainty one would have about the composition of a subgroup chosen at random. Thus, we can use the entropy of subgroup composition to compare different species or situations. Moreover, this degree of uncertainty may be a relevant feature not only for the researchers but for the animals themselves. Intuitively,

66 individuals in a group with a high degree of fission-fusion dynamics would face more uncertainty
67 about the composition of the subgroups they can form than individuals of species with less
68 flexible grouping patterns. The more uncertainty in the identity of group-mates, the greater the
69 uncertainty in social interactions [14]. Dealing with such uncertainty is thought to present a
70 cognitive challenge [7, 12, 15].

71 We thus propose that Shannon's entropy can be used to quantify social uncertainty due to fission-
72 fusion dynamics at the group and individual levels. At the group level, Shannon's entropy has
73 been used for characterising the overall degree of variation and uncertainty in social networks
74 [14, 16, 17]. Accordingly, we propose that the entropy of subgroup composition can be used as a
75 general metric of this particular dimension of fission-fusion dynamics [1]. At the individual level,
76 Shannon's entropy could also reflect the uncertainty actually faced by individuals in these groups.
77 Shannon's entropy has been used to quantify how evenly an individual distributes its grooming
78 interactions amongst the rest of the individuals in its group [8]. Our proposal is analogous
79 to this use of Shannon's entropy, but applies to the spatiotemporal associations between an
80 individual and the rest of its group mates. When subgroup composition is highly variable,
81 individuals do not repeat their interactions with the same individuals often. A lower frequency
82 of repeated interactions may lead to a higher uncertainty about social relationships, which in
83 turn may require alternate ways of reducing such uncertainty and predicting the outcome of
84 social interactions and others' behavior [7, 14]. Our approach to quantifying this uncertainty
85 should be relevant to any species exhibiting some degree of fission-fusion dynamics, where group
86 members repeat interactions with others, finding themselves associated with others at different
87 frequencies and individually recognizing one another (or at least classifying other group members
88 in broad categories) [1, 18].

89 We develop a proof of concept by measuring Shannon's entropies at the group and individual
90 levels in three species that show different degrees of fission-fusion dynamics: spider monkeys
91 (*Ateles geoffroyi*), chimpanzees (*Pan troglodytes*) and gelada monkeys (or geladas, *Theropithecus*
92 *gelada*), although our approach should be applicable to any species where the composition of
93 subgroups can be reliably observed and quantified. Although these three primates are known
94 for their high variability in subgroup size and cohesion, they differ in the degree of variation

in subgroup composition. Spider monkey and chimpanzee subgroups are highly variable in composition with group members fissioning and fusing independently from one another [19]. In contrast, geladas have a multi-level social system with highly stable one-male units that fission and fuse with one another in predictable ways, creating a higher order “band” structure [20–22]. Because of this, we predict geladas to have lower entropy values than spider monkeys and chimpanzees, despite the fact that they live in larger groups. We quantify social uncertainty at the group and individual levels using Shannon’s entropy and a randomized expectation of entropy that considers subgroup size variation and sample size, against which the observed entropy can be compared. We complement the estimation of social uncertainty with an estimation of the number of subgroups that the group is likely to be divided into at any given time, based on the observed composition and size of single focal subgroups.

2 Methods

2.1 Data collection

Spider monkey data were collected from August 2009 to July 2010 and from January 2013 to September 2014 in the *Otoch Ma’ax yetel Kooh* protected area, in the Yucatan peninsula, Mexico. The study group has been monitored continuously since 1997, and all group members are identified and habituated to human presence. Observations consisted of instantaneous scan samples performed every 20 minutes between 0600 and 1800 hours on subgroups chosen according to criteria for homogenizing sample size across individuals. A total of 3916 scan samples, equivalent to 1305 h of observation, were collected in the 2009-2010 period and a total of 7917 scan samples, equivalent to 2639 h of observation, were collected in the 2013-2014 period. During each scan sample, the identities of all subgroup members were recorded. A subgroup was defined using a chain rule of 30m, such that individuals 30m or closer to any other were considered as part of the same subgroup [23, 24]. Only adult individuals were included in the analysis: 10 females and 7 males in 2009-2010 and 18 females and 7 males in 2013-2014.

Chimpanzee data were collected from January 2008 to December 2009 from the Sonso commu-

nity in the Budongo Forest, Uganda. The study group has been monitored continuously since 1990 [25]. All group members are identified and habituated to human presence. Observations consisted of instantaneous scan samples performed every 15 minutes during focal follows between 0600 and 1800 hours, recording the identities of all subgroup members. A subgroup was defined as all individuals visible or known to be present within 35-50m of the focal animal [25]. Subgroups were chosen each day according to criteria for homogenizing sample size across individuals. The 2008 period contained a total of 10616 scan samples, equivalent to 2654 h of observation, while the 2009 period contained a total of 12935 scan samples, equivalent to 3234 h of observation. Only adult individuals that were present throughout each of the two entire years were included in the analysis: a total of 20 (in 2008) and 21 (in 2009) females and 9 males (both years).

Gelada data were collected from January 2014 to December 2015 in a population that has been continuously monitored since 2006 in the Simien Mountains National Park, Ethiopia [22]. Each morning, observers recorded the identity of all known individuals in a gelada subgroup (defined using a chain rule of 50m) and then followed it for 1-8 hours. During follows, the observers collected a scan sample every 30 minutes, recording the identity of all known individuals currently in the subgroup. The 2014 period consisted of 1420 scan samples, equivalent to 473 h of observation, and the 2015 period consisted of a total of 1168 scan samples, equivalent to 389 h of observation. Only adult individuals were included in the analysis: 21 males and 82 females in 2014 and 29 males and 97 females in 2015.

2.2 Entropy calculation

To quantify social uncertainty at the group level, we calculated Shannon’s entropy of subgroup composition as follows. Imagine that a large number of observations allows the accurate estimation of the probability of occurrence of any particular subgroup (or subset) with composition $\{a\}$:

$$p_{\{a\}} \equiv \frac{\text{number of observed subsets} = \{a\}}{\text{total number of observed subsets}} \tag{2}$$

146 in a group (set) of n elements. The composition entropy H of the group stems from the definition
 147 (1):

$$H = - \sum_{\{a\}} p_{\{a\}} \log_2 p_{\{a\}}, \quad (3)$$

148 where the sum runs over all the observed compositions, i.e. those with $p_{\{a\}} \neq 0$.

149 To quantify social uncertainty at the individual level, we applied a similar entropy formula, but
 150 from the perspective of each individual. For those subgroups in which a given individual i was
 151 present, we measured i 's entropy by considering the different compositions of the subgroup in
 152 terms of the remaining $n - 1$ individuals (see Figure 1).

153 For each entropy, it is useful to determine an upper bound value, denoted as H^* , through a
 154 null model neglecting preferential associations: The subsets of size k appear with the same
 155 size frequency distribution f_k as in observations, where $\sum_{k=1}^n f_k = 1$ by normalization, but all
 156 the compositions of same size k are assumed to be equiprobable. Given a particular subgroup
 157 composition $\{a\}$ of k_a individuals, there are $C_{k_a}^n$ different ways of choosing k_a elements from
 158 n , where $C_k^n = n! / [(n - k)!k!]$ stands for “ n choose k ”. The null conditional probability of $\{a\}$
 159 given k_a , $p^*(\{a\}|k_a)$, takes the form $p^*(\{a\}|k_a) = 1/C_{k_a}^n$. One deduces:

$$p_{\{a\}}^* = p^*(\{a\}|k_a) f_{k_a} = \frac{f_{k_a}}{C_{k_a}^n}, \quad (4)$$

160 for the null composition probability $p_{\{a\}}^*$. The null maximal entropy follows:

$$H^* = - \sum_{\{a\}} p_{\{a\}}^* \log_2 p_{\{a\}}^* = - \sum_{k=1}^n f_k \ln_2 \left(\frac{f_k}{C_k^n} \right), \quad (5)$$

161 where, in the last equality, one has used the fact that in the sum over all compositions the terms
 162 can be re-arranged by size: each size k as a fixed factor $p_{\{a\}}^* \ln_2 p_{\{a\}}^*$, which appears C_k^n times
 163 in the sum.

2.3 Bootstrap entropy

The number of observations being finite in any empirical data set, it is often problematic to evaluate the probabilities $p_{\{a\}}$ by using Eq. (2), since many compositions of low probability may not be observed and are thus replaced by zero in the sum (3). Therefore the empirical H resulting from N_o observations *a priori* underestimates the real entropy. For a fair comparison of H with a randomized model, it is thus necessary to calculate the entropy of the randomized model given N_o observations as well, instead of Eq. (5). This can be done numerically with a bootstrap, or analytically as follows.

Let us denote $N(k) = N_o f_k$ as the number of times subgroups of size k have been observed in the data, with $\sum_{k=1}^n N(k) = N_o$. Let us denote $n_{\{a\}}$ as the number of times a given composition $\{a\}$ (of size k_a) is observed from a sampling of size N_o of the null model. The probability that $\{a\}$ appears exactly i times [$i = 0, \dots, N(k_a)$] in this sampling is given by the binomial distribution:

$$\text{Prob}[n_{\{a\}} = i] = p^*(\{a\}|k_a)^i [1 - p^*(\{a\}|k_a)]^{N(k_a)-i} C_i^{N(k_a)}, \quad (6)$$

where $p^*(\{a\}|k_a) = 1/C_{k_a}^n$. The bootstrap entropy H^b is obtained by replacing $p_{\{a\}}$ by $n_{\{a\}}/N_o$ in Eq. (3) and taking the average over all the possible values of $n_{\{a\}}$:

$$H^b = - \sum_{\{a\}} \left\langle \frac{n_{\{a\}}}{N_o} \ln_2 \left(\frac{n_{\{a\}}}{N_o} \right) \right\rangle = - \sum_{\{a\}} \sum_{i=1}^{N(k_a)} \frac{i}{N_o} \ln_2 \left(\frac{i}{N_o} \right) \times \text{Prob}[n_{\{a\}} = i]. \quad (7)$$

(The term $i = 0$ contributes to 0.) Making the substitution $\sum_{\{a\}} \rightarrow \sum_{k=1}^n C_k^n$ as in (5) and using Eq. (6), one obtains the bootstrap entropy:

$$H^b = - \sum_{k=1}^n C_k^n \sum_{i=1}^{N(k)} \frac{1}{[C_k^n]^i} \left[1 - \frac{1}{C_k^n} \right]^{N(k)-i} C_i^{N(k)} \frac{i}{N_o} \ln_2 \left(\frac{i}{N_o} \right). \quad (8)$$

One recovers Eq. (5) by taking the limit $N_o \rightarrow \infty$ and $N(k) \rightarrow \infty$, keeping $N(k)/N_o = f_k$ fixed. This entropy is a more useful point of comparison with the observed data, because in contrast with the null maximal entropy H^* , where all combinations are equally likely regardless of the sample size, H^b is computed with the sample size of the observed data, the composition of the

subgroups being randomised. It is also equal to the mean entropy for a set of bootstrapped original data, in which the 1s and 0s in Figure 1 have been randomly shuffled within rows, thus keeping subgroup size and the number of observations for each individual unchanged.

2.4 Entropy comparisons

The Kullback-Leibler (KL) divergence is commonly used to quantify how much an empirical distribution $p_{\{a\}}$ differs from an expectation $p_{\{a\}}^*$, thus providing a way of comparing the observed entropy H to the null maximal entropy H^* [26]. It is defined as $KL(p|p^*) = \sum_{\{a\}} p_{\{a\}} \ln_2(p_{\{a\}}/p_{\{a\}}^*)$ and represents, in the present context, the average additional amount of information (in bits) needed to identify a subgroup randomly drawn from $p_{\{a\}}$, when assuming that the distribution is $p_{\{a\}}^*$. For the null maximal model, it reads:

$$KL(p|p^*) = \sum_{\{a\}} p_{\{a\}} \ln_2 \left(\frac{p_{\{a\}} C_{k_a}^n}{f_{k_a}} \right), \quad (9)$$

where, once again, k_a represents the size of $\{a\}$, $p_{\{a\}}$ is given by Eq. (2), and the sum is over observed compositions.

The above quantity cannot be applied to compare H to the bootstrap entropy H^b , however, since a finite sample of $p_{\{a\}}^*$ can contain vanishing entries, thus making KL infinite. A useful alternate measure is the Jensen-Shannon distance between two distributions p and q , defined by $J(p|q) = H(m) - \frac{1}{2}[H(p) + H(q)]$, where the entries of m are $m_{\{a\}} = (p_{\{a\}} + q_{\{a\}})/2$ [26]. Unlike the KL divergence, J can deal with vanishing entries (i.e., $\{a\}$ such that $q_{\{a\}} = 0$), it is symmetric and satisfies the triangle inequality. Another interesting property is that J is 0 if $p = q$ and 1 if the distributions have completely disjoint supports. This distance is thus adimensional, making comparisons across systems easier *a priori*. Here, q is a sampling of size N_o of the null model, and we calculate the average of $J(p|q)$, denoted as J^b , over all possible samplings. Following the same steps leading to Eq. (8), the mean distance between observations

206 and the bootstrap reads:

$$J^b = 1 - \frac{1}{2} \sum_{\{a\}} \sum_{i=1}^{N(k_a)} \frac{1}{[C_{k_a}^n]^i} \left[1 - \frac{1}{C_{k_a}^n} \right]^{N(k_a)-i} C_i^{N(k_a)} \left[p_{\{a\}} \ln_2 \left(1 + \frac{i}{N_o p_{\{a\}}} \right) + \frac{i}{N_o} \ln_2 \left(1 + \frac{N_o p_{\{a\}}}{i} \right) \right]. \quad (10)$$

207 where the sum runs over observed compositions.

208 The methods for the partition analysis are included in the ESM. All analyses were implemented
209 in R [27] and the code is shared in the ESM.

210 **3 Results**

211 **3.1 Social uncertainty at the group level**

212 The values of entropy (H) at the group level were highly consistent: 8 - 8.5 for spider monkeys,
213 close to 10 for chimpanzees and close to 5 for geladas (Figure 2) and significantly lower than
214 the calculated and bootstrap entropies (H^b) in each case; which is confirmed by KL divergences
215 and JS distances (Figure 2), which were especially large for geladas.

216 When considering male-only data for geladas, entropy was relatively unchanged (2014: observed
217 5.76 bits, bootstrap 10.12 bits; 2015: observed 6.19 bits, bootstrap 10.16 bits) but the KL
218 divergence was lower (2014: KL divergence 9.67; 2015: 16.9; compare to values in Figure 2,
219 which are around 70 and 90, for 2014 and 2015, respectively). On the contrary, the JS distance
220 is still close to the maximal value of 1 when considering only the males (2014: 0.95; 2015: 0.99;
221 compare to similar values in Figure 2).

222 **3.2 Social uncertainty at the individual level**

223 Figure 3 shows summaries of the entropy from the perspective of different individuals. For spider
224 monkeys (Figure 3a-b), individual entropy varied from 4 to 8 bits in the case of females and
225 tends to be lower and more consistent in the case of males. While the majority of individuals in

2009 show H values that are clearly lower than the bootstrap entropy H^b values, in 2013 there are several females that show H values that are very close to H^b . When comparing these values using the JS distance (Figure S4), there are indeed some females in the 2013 dataset for whom the JS distance between the observed and the bootstrap entropy is relatively low (< 0.85), and who could be considered to show a particularly high entropy compared to other females. Their subgroups, thus, seem to show a maximum level of variability in composition.

In the case of chimpanzees, we found an opposite effect of sex on the individual entropy values: males tend to have a higher and less variable observed entropy than females (Figure 3c-d). The H values for females ranged from 0.88-9.56 bits and those of males ranged from 8.2-9.9 bits. In all individuals, H was clearly lower than H^b in both periods, as can be confirmed by the JS distance values (Figure S5). The values of JS distance (Figure S5) show more variability amongst individual females than amongst the males.

Gelada individual entropy is aligned to the one-male unit to which individuals belong (Figure 3e-f), resulting in females sharing the same individual entropy values as the male. As in the case of the group entropy values (Figure 2), individual entropy values were farther from the bootstrap expectation than in the case of the other two species. Comparing these values using the JS distance is not very useful, as most values lie very close to 1 (data not shown). However, limiting the analysis to the one-male units yielded variability in the JS distances, particularly in the 2013 dataset (Figure S6). Here, some one-male units have a JS that is farther away from the maximum number, indicating that they have a higher degree of variability in their subgroup compositions.

The results of the partition analysis, aimed at establishing the probability that a group is split into different numbers of subgroups, can be found in the Electronic Supplementary Material.

249 4 Discussion

We used Shannon's entropy to quantify temporal variation in subgroup composition across three primate species and provide a measure of "social uncertainty" at the group and individual levels. As predicted, spider monkeys and chimpanzees, considered as species with a high degree

253 of fission-fusion dynamics [1, 19], have a higher entropy of subgroup composition than geladas,
254 which show variation in subgroup size and spatial cohesion between group members, but also
255 have indivisible one-male units and a higher order band structure [20]. This latter characteristic
256 constrains the flexibility in subgroup composition of geladas compared to spider monkeys and
257 chimpanzees, and explains why geladas have a lower observed entropy.

258 That variation in sample size and group size within a species does not influence the estimation
259 of entropy suggests that our method is robust and could be used to compare social uncertainty
260 across different species and datasets with different characteristics. The bootstrap entropy H^b ,
261 corresponding to the maximum entropy that could be expected in a dataset of the same size and
262 subgroup size distribution if all subgroup compositions were equally likely, serves as a reference
263 to evaluate whether the observed entropy is relatively high or low. In all three species, that
264 the observed entropy is lower than the bootstrap entropy implies that preferential associations
265 between individuals make some compositions more likely than others among the full set of
266 potential compositions. Together, the observed and bootstrap entropies serve as a measure of
267 how much of the potential variation in subgroup composition is actually observed.

268 We propose that our method can be used to compare the degree of fission-fusion dynamics
269 between species, particularly in terms of the temporal variation in subgroup composition. In
270 particular, Jensen-Shannon (JS) distances serve as a quantification of how far the observed
271 entropy is from the bootstrap entropy and can be used as for comparative purposes. A species
272 with a high JS distance (close to unity) would have a variation in subgroup composition that is
273 far from the maximum expected by the bootstrap entropy, and would have a relatively low degree
274 of temporal variation in subgroup composition. Conversely, a species with a low JS distance
275 would have a variation in subgroup composition that is close to the maximum expected, and
276 thus would have a relatively high degree of temporal variation in dimension of fission-fusion
277 dynamics. In our analysis, geladas stood out as having the highest JS distance relative to the
278 bootstrap entropy and thus would be the species with the lowest degree of variation in subgroup
279 composition. The implication is that there are more constraints to the flexibility of association,
280 and thus a lower uncertainty in subgroup composition, in geladas than in the other two species.
281 The difference in JS distances between spider monkeys and chimpanzees, although not as high

282 as between these two species and the geladas, is still detectable and suggests that chimpanzees
283 have the highest degree of temporal variation in this dimension of fission-fusion dynamics of the
284 three analyzed species. JS distances can also be used to explore within-species differences in the
285 temporal variation in subgroup composition. As Figure 3 shows, for spider monkeys JS distance
286 is larger in 2009 than in 2013, whereas JS distances of the two chimpanzee data sets are rather
287 similar.

288 The level of analysis of social uncertainty for geladas deserves special attention. The existence
289 of indivisible, one-male units decreases the number of potential subgroup compositions and thus
290 the observed entropy when considering all individuals. We also ran the analysis considering
291 only males, thus estimating the degree of flexibility of association between one-male units.
292 We obtained similar values of entropy at both levels, with an observed entropy around 4 bits
293 lower than the bootstrap expectation. This result is what would be expected if one-male units
294 associated preferentially with a few of the other one-male units, as opposed to associating freely
295 with all units. In other words, a low value of entropy in the association of one-male units into
296 bands (i.e. the clearest, more consistently observed level above the one-male units; [22]) implies
297 that the composition of these bands is relatively predictable. The JS distances between observed
298 and bootstrap entropies when considering only males were close to 1, suggesting that there is
299 much less variation than could be expected if there were no preferential associations between
300 one-male units. However, the fact that there are many more males in 2014 apparently leads to
301 much more predictable patterns (i.e. JS distances close to unity in all cases). It is as if the one-
302 male units responded by becoming less fluid, perhaps as a way of maintaining a low uncertainty
303 in the face of an increase in group size and all the potential disorder (i.e. higher entropy)
304 this could cause. The nonrandom association of one-male units in this population of geladas
305 has been demonstrated using both social network analysis [21] and hierarchical clustering [22].
306 Thus, our results are consistent with what we know about gelada multilevel society, but they
307 go a step further by quantifying a component of social complexity that is closely related to
308 social uncertainty due to fission-fusion dynamics and that can be compared between and within
309 species.

310 We also extended our analysis to the entropy of subgroup composition from the point of view

of each group member. Overall, we obtained values similar to those for the whole group, but some differences between individual values of entropy were revealing. In spider monkeys and chimpanzees, the two sexes differed in their individual entropy values. Higher values in female spider monkeys compared to males are consistent with the females' lower rates of preferential association compared to males [28]. By contrast, lower values in female chimpanzees compared to males could be due to their known tendency to form strong and lasting bonds with particular females [29], as well as to the opportunistic nature of associations between males [30]. In spider monkeys, the females with a lowest JS distance had recently immigrated into the group (i.e. females AE, HI, PC and TG in Figure S4b). This result suggests that is consistent with previous studies [28] that found that during their first year in the group, immigrant females' preference for particular others is low. This is an example of the utility of comparing the observed and bootstrap entropy values using the JS distance. In the case of chimpanzees, the 2009 data contained a newly immigrating female (TJ), which also had a relatively high JS distance value compared to other individuals (Figure 5Sb). Other females with particularly high JS distances include BC, KG and FL, who had severe snare injuries (entire hand or foot missing), and thus limited their movements to the core area of the home range and were observed in smaller subgroups than the rest of the females. This is an example of the usefulness of comparing observed entropy values between individuals.

An individual's entropy value can be interpreted as the degree of uncertainty it has about its particular set of associations [14]. It has long been established that several social interactions are aimed at reducing the stress caused by uncertainty in social relationships [31,32]. A reduction in uncertainty has been proposed to lie at the core of emerging features of social structure such as dominance hierarchies [14,33,34]. In species where repeated social interactions occur amongst group members that form subgroups, our measures of entropy at the individual level are a promising metric for quantifying social uncertainty due to fission-fusion dynamics and for comparing this component of social complexity across individuals, situations, groups and species. Individuals with a lower observed entropy relative to the bootstrap entropy would face less uncertainty than individuals with similar values of observed and bootstrap entropy. Analysing these individual differences may help researchers understand the role played by individuals in

340 their groups and the extent to which they could predict the interactions amongst others in the
341 group [35].

342 One of the reasons a high degree of fission-fusion dynamics is considered to be cognitively chal-
343 lenging is that individuals face a high uncertainty about their social relationships [1]. For social
344 interactions to reduce the uncertainty about other group member's behavior and the quality of
345 relationships with them [31], specific mechanisms must be in place that can allow individuals to
346 update their information about these relationships with others, as well as to generalize across
347 different relationships that share similar features. Therefore, cognitive abilities that allow in-
348 dividuals to reduce their uncertainty with respect to social relationships, like abstraction (e.g.
349 using concepts such as "friend" or "potential mate" to classify relationships) and transitivity
350 (i.e. inferring a linear order of relationships using partial information), may be particularly
351 important in species with high levels of fission-fusion dynamics, where the understanding of
352 social relationships must be carried out using partial information in highly variable social con-
353 texts [6]. In addition, cognitive abilities to deal with uncertainty, such as inhibition of ongoing
354 responses until the social situation can be assessed when subgroup composition changes, are
355 also important in fission-fusion dynamics [36]. We predict that species with a high uncertainty
356 in subgroup composition are more likely to show these cognitive abilities than species with a
357 lower uncertainty.

358 Estimating the probability that the whole group would be partitioned, or split, in different
359 numbers of subgroups provides a further way to quantify social uncertainty. The probability
360 distributions that result from our partition analysis can be considered a measure of the uncer-
361 tainty with respect to the grouping patterns of unobserved group members. For example, it
362 might be easier for an individual to predict which group members not present in its current
363 subgroup could be close or associated with one another in a group that is potentially split in
364 2-6 subgroups than in a group that is split in 9-14 subgroups (e.g. compare Figures S9 a and
365 b). In addition to its usefulness for studying higher levels in multi-level societies, our partition
366 analysis could be more generally applied in any study in which only one subgroup can be fol-
367 lowed at one time (like in the majority of studies of species with a high degree of fission-fusion
368 dynamics). For example, research on topics like between-subgroup vocal interactions [23,37] or

home ranges [38,39] could be aided by an estimation of how many subgroups there are likely to be at a given time, even if only one subgroup has been monitored directly.

It is necessary to note that our method assumes that the distribution of observed subgroup size f_k reflects the true distribution of subgroup size in which a group was found during a certain study period. Under that assumption, the bootstrap entropy H^b reflects the maximum entropy that could be observed given the observed distribution of subgroup size. Also, our estimation of the most likely partition in which the group is found relies on a correctly estimated f_k . However, when studying species with high degrees of fission-fusion dynamics, there are potential biases which might make it more likely for researchers to observe the larger or more conspicuous subgroups. Thus, in field studies, steps should be taken to ensure that the sample of subgroups is representative of the true distribution.

Establishing metrics to estimate social complexity is not a trivial matter [12,40,41]. Crude measures, such as group size, number of different interactions, presence of triadic interactions, etc., have been used but have not been operationalized in such a way that different species with different group size and degree of fission-fusion dynamics can be compared (but see [10]). As we show, Shannon's entropy represents a relevant metric of social uncertainty as one component of social complexity, but it is important to bear in mind the relationship between complexity and uncertainty. While a completely random process, which in turn would have the highest entropy, would be maximally uncertain, we would not necessarily consider it as a complex process. On the opposite end, a fully predictable pattern, with minimal complexity, would also be minimally uncertain, with a correspondingly low entropy. When considering complexity, including social complexity, we need to take into account both the flexibility and the nonrandom structure of a process [42, p. 353] [43]. Thus, maximally complex societies would not necessarily lie in any of the two extremes of the uncertainty spectrum. A middle-ground, where relationships are somewhat predictable, also corresponds to the greatest degree of relationship differentiation, which is another way to characterize social complexity [10]. This is because random processes would involve no relationship differentiation, while completely stable groups can emerge from simple rules that involve only a categorical differentiation between in and out-group individuals. We predict that, in terms of subgroup composition, higher social complexity would occur in

398 groups with high observed entropy that is, nonetheless, still lower than the bootstrap entropy.
399 In terms of JS distances, a species would have a higher social complexity at intermediate values.
400 In these groups, individuals would have to cope with a high degree of uncertainty about who
401 their associates would be at any one time, but at the same time maintain a diversity of social
402 relationships with preferred companions, in many different contexts [10,11,32,34]. It is possible
403 that the real complexity might lie in the cognitive and behavioral mechanisms used to deal with
404 social uncertainty in the face of an existing social structure.

405 Our approach to measuring social complexity through social uncertainty can be applied to
406 any species that interacts in temporary and variable subsets and may be particularly relevant
407 for taxa in which a known set of individuals can recognize one another through visual, vocal
408 or olfactory means. The proposed metrics should also be useful for future studies comparing
409 the degree of fission-fusion dynamics across species varying to different extent in subgroup
410 composition, together with subgroup size and spatial cohesion [1]. More generally, they can aid
411 our understanding of the influence of flexible social settings on the interactions between group
412 members and their implications for social cognition.

413 **5 Ethics**

414 All fieldwork was conducted under the Guidelines for the Use of Animals in Research of the
415 Association for the Study of Animal Behaviour / Animal Behavior Society and conformed to
416 the legal requirements of the respective countries where it was conducted.

417 **6 Data and code**

418 All datasets used to illustrate the methodology are shared in a data repository. We also share
419 the code in R [27] to calculate the entropy measures, both at the group and individual levels, the
420 comparison of entropies using the KL divergences and the JS distances, as well as the partition
421 analysis (see Electronic Supplementary Material).

422 **7 Competing interests**

423 We have no competing interests.

424 **8 Authors' contributions**

425 The study was conceived by GRF, AJK, FA and DB. The study was designed by GRF, AJK,
426 MCC, AD, JL, CMS, FA and DB. Field data were collated by JL, JCB, TCB, NSM and KS.
427 Data analysis was carried out by GRF and DB, with techniques developed by DB. The workshop
428 where ideas were first discussed was coordinated by FA and CMS. The manuscript was drafted
429 by all coauthors, who also gave final approval for publication.

430 **9 Funding**

431 This study was first discussed at a Workshop on Fission-Fusion Dynamics funded by the Wenner-
432 Gren Foundation (FA and CMS). Funding was provided by CONACYT and Instituto Politécnico
433 Nacional (GRF), Santander Mobility Award/Swansea University (AJK), Wildlife Conservation
434 Society, National Geographic Society, Leakey Foundation, National Science Foundation and the
435 National Institute on Aging (TJB, JCB and NSM), National Geographic and Chester Zoo (FA
436 and CS), PAPIIT-UNAM 105015 (DB), NSF IOS-1255974 (TB and JB) and NSF III 1514174
437 and David and Lucile Packard Foundation 2016-65130 (MCC).

438 **10 Acknowledgements**

439 FA, CS and GRF thank the field assistants at the Yucatan spider monkey field site (Augusto,
440 Eulogio, Juan and Macedonio Canul) and Braulio Pinacho-Guendulain and Sandra E. Smith
441 Aguilar for collecting and collating data on spider monkeys. Vernon Reynolds shared data from
442 the long-term field study on the Budongo chimpanzees. TB, JB and NSM extend their thanks to
443 members of the Simien Mountains Gelada Research Project for their tireless help in the field and

the Ethiopian Wildlife Conservation Authority for their permission to work with the geladas. AJK thanks Ines Fürtbauer for discussion. A Santander Mobility Award/Swansea University funds enabled collaboration between AJK, GRF and DB during the writing of this paper. DB thanks the Max Planck Institute for the Physics of Complex Systems for hospitality. We thank two anonymous reviewers for their helpful comments.

References

- [1] Aureli F, Schaffner CM, Boesch C, Bearder SK, Call J, Chapman CA, et al. Fission-fusion dynamics: new research frameworks. *Current Anthropology*. 2008;49(4):627–654.
- [2] Kummer H. *Primate societies: Group techniques of ecological adaptation*. Chicago: Aldine-Atherton; 1971.
- [3] Chapman CA, Chapman LJ, Wrangham R. Ecological constraints on group size: an analysis of spider monkey and chimpanzee subgroups. *Behavioral Ecology and Sociobiology*. 1995;36(1):59–70.
- [4] Smith JE, Kolowski JM, Graham KE, Dawes SE, Holekamp KE. Social and ecological determinants of fission–fusion dynamics in the spotted hyaena. *Animal Behaviour*. 2008;76(3):619–636.
- [5] Asensio N, Korstjens AH, Aureli F. Fissioning minimizes ranging costs in spider monkeys: a multiple-level approach. *Behavioral Ecology and Sociobiology*. 2009;63(5):649–659.
- [6] Cheney DL, Seyfarth RM. *How monkeys see the world: Inside the mind of another species*. Chicago, IL: Chicago University Press; 1990.
- [7] Bergman TJ, Sheehan MJ. Social knowledge and signals in primates. *American journal of primatology*. 2013;75(7):683–694.
- [8] Cheney DL. Intragroup cohesion and intergroup hostility: the relation between grooming distributions and intergroup competition among female primates. *Behavioral Ecology*. 1992;3(4):334–345.

- 469 [9] DiBitetti MS Mario SBitetti. The distribution of grooming among female primates: testing
470 hypotheses with the Shannon-Wiener diversity index. *Behaviour*. 2000;137(11):1517–1540.
- 471 [10] Fischer J, Farnworth MS, Sennhenn-Reulen H, Hammerschmidt K. Quantifying social
472 complexity. *Animal Behaviour*. 2017;130:57–66.
- 473 [11] Freeberg TM, Dunbar RI, Ord TJ. Social complexity as a proximate and ultimate factor
474 in communicative complexity. *Philosophical Transactions of the Royal Society of London*
475 *B: Biological Sciences*. 2012;367(1597):1785–1801.
- 476 [12] Bergman TJ, Beehner JC. Measuring social complexity. *Animal Behaviour*. 2015;103:203–
477 209.
- 478 [13] Shannon C. A mathematical theory of communication. *Bell Systems Technical Journal*.
479 1948;27:379–423.
- 480 [14] Barrett L, Henzi SP, Lusseau D. Taking sociality seriously: the structure of multi-
481 dimensional social networks as a source of information for individuals. *Philosophical Trans-*
482 *actions of the Royal Society of London B: Biological Sciences*. 2012;367(1599):2108–2118.
- 483 [15] Dunbar RI, Shultz S. Bondedness and sociality. *Behaviour*. 2010;147(7):775–803.
- 484 [16] Bailey KD. Social entropy theory. SUNY Press; 1990.
- 485 [17] Haddadi H, King AJ, Wills AP, Fay D, Lowe J, Morton AJ, et al. Determining associ-
486 ation networks in social animals: choosing spatial temporal criteria and sampling rates.
487 *Behavioral Ecology and Sociobiology*. 2011;65(8):1659–1668.
- 488 [18] Loretto MC, Schuster R, Itty C, Marchand P, Genero F, Bugnyar T. Fission-fusion dy-
489 namics over large distances in raven non-breeders. *Scientific Reports*. 2017;7(1):380.
- 490 [19] Symington MM. Fission-fusion social organization in *Ateles* and *Pan*. *International Journal*
491 *of Primatology*. 1990;11(1):47–61.
- 492 [20] Dunbar R. Social dynamics of gelada baboons. Switzerland: Basel: Karger; 1975.
- 493 [21] Mac Carron P, Dunbar R. Identifying natural grouping structure in gelada baboons: a
494 network approach. *Animal Behaviour*. 2016;114:119–128.

- [22] Snyder-Mackler N, Beehner JC, Bergman TJ. Defining higher levels in the multi-level societies of geladas (*Theropithecus gelada*). *International Journal of Primatology*. 2012;33(5):1054–1068.
- [23] Ramos-Fernández G. Vocal Communication in a Fission-Fusion Society: Do Spider Monkeys Stay in Touch With Close Associates? *International Journal of Primatology*. 2005 Oct;26(5):1077–1092.
- [24] Aureli F, Schaffner CM, Asensio N, Lusseau D. What is a subgroup? How socioecological factors influence interindividual distance. *Behavioral Ecology*. 2012;23(6):1308–1315.
- [25] Reynolds V. The chimpanzees of the Budongo forest: Ecology, behaviour and conservation. Oxford University Press; 2005.
- [26] Lin J. Divergence measures based on the Shannon entropy. *IEEE Transactions on Information theory*. 1991;37(1):145–151.
- [27] R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing; 2016.
- [28] Ramos-Fernandez G, Boyer D, Aureli F, Vick LG. Association networks in spider monkeys (*Ateles geoffroyi*). *Behavioral Ecology and Sociobiology*. 2009;63(7):999–1013.
- [29] Foerster S, McLellan K, Schroepfer-Walker K, Murray CM, Krupenye C, Gilby IC, et al. Social bonds in the dispersing sex: partner preferences among adult female chimpanzees. *Animal behaviour*. 2015;105:139–152.
- [30] Gilby IC, Wrangham RW. Association patterns among wild chimpanzees (*Pan troglodytes schweinfurthii*) reflect sex differences in cooperation. *Behavioral Ecology and Sociobiology*. 2008;62(11):1831.
- [31] Van Hooff J, Aureli F. Social homeostasis and the regulation of emotion. *Emotions: Essays on emotion theory*. 1994;p. 197–217.
- [32] Wittig RM, Crockford C, Lehmann J, Whitten PL, Seyfarth RM, Cheney DL. Focused grooming networks and stress alleviation in wild female baboons. *Hormones and Behavior*. 2008;54(1):170–177.

- [33] Drews C. The concept and definition of dominance in animal behaviour. *Behaviour*. 1993;125(3):283–313.
- [34] Flack JC. Multiple time-scales and the developmental dynamics of social systems. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*. 2012;367(1597):1802–1810.
- [35] Flack J. 12 Life's Information Hierarchy. *From Matter to Life: Information and Causality*. 2017;p. 283.
- [36] Amici F, Aureli F, Call J. Fission-fusion dynamics, behavioral flexibility, and inhibitory control in primates. *Current Biology*. 2008;18(18):1415–1419.
- [37] Cantor M, Shoemaker LG, Cabral RB, Flores CO, Varga M, Whitehead H. Multilevel animal societies can emerge from cultural transmission. *Nature communications*. 2015;6:8091.
- [38] Smith-Aguilar SE, Ramos-Fernández G, Getz WM. Seasonal changes in socio-spatial structure in a group of free-living spider monkeys (*Ateles geoffroyi*). *PloS one*. 2016;11(6):e0157228.
- [39] Sprogis KR, Raudino HC, Rankin R, MacLeod CD, Bejder L. Home range size of adult Indo-Pacific bottlenose dolphins (*Tursiops aduncus*) in a coastal and estuarine system is habitat and sex-specific. *Marine Mammal Science*. 2016;32(1):287–308.
- [40] Whiten A. Social complexity and social intelligence. *The nature of intelligence*. 2000;p. 185–196.
- [41] Whitehead H. Analyzing animal societies: quantitative methods for vertebrate social analysis. Chicago, IL: University of Chicago Press; 2008.
- [42] Gleick J. *The Information: A History, A Theory, A Flood*. Vintage Books, NY; 2012.
- [43] Prokopenko M, Boschetti F, Ryan AJ. An information-theoretic primer on complexity, self-organization, and emergence. *Complexity*. 2008;15(1):11–28.

Individuals							Group entropy: observed subgroup compositions	Individual entropy: observed compositions for individual A
A	B	C	D	E	F	G		
●	●	●	●	○	○	○	1 1 1 1 0 0 0	1 1 1 0 0 0
●	●	●	●	○	○	○	1 1 1 1 0 0 0	1 1 1 0 0 0
●	●	○	○	○	○	○	1 1 0 0 0 0 0	1 0 0 0 0 0
●	●	○	○	○	○	○	1 1 0 0 0 0 0	1 0 0 0 0 0
●	●	○	○	●	●	●	1 1 0 0 1 1 1	1 0 0 1 1 1
○	●	○	○	●	●	●	0 1 0 0 1 1 1	

Figure 1: Dataset coding for the calculation of subgroup entropy, at group and individual levels. The data consist of observations at regular intervals (rows) on different individuals who can be present (filled circles) or absent (empty circles) in any given subgroup due to fissions and fusions. For calculating the group entropy, we code presences as 1 and absences as 0 and each subgroup composition would correspond to a particular sequence of 1 and 0. For calculating the individual entropy for an individual A, we do the same but only for those subgroups in which A was present and considering all other individuals except A (shaded area), thus capturing the variability in subgroup composition from A's perspective.




Figure 2: (a) entropy of subgroup composition for two data sets of each of three species: spider monkeys (blue; data from 2009 and 2013), chimpanzees (green; data from 2008 and 2009) and geladas (dark red; data from 2014 and 2015). Solid dots correspond to the observed entropy (H) and empty dots to the bootstrap entropy (H^b). Group size n is 17 and 25 for spider monkeys, 29 and 30 for chimpanzees and 103 and 126 for geladas. Sample size No is 3916 and 7917 for spider monkeys, 10616 and 12935 for chimpanzees and 1420 and 1168 for geladas. (b) KL divergence between the observed and the null maximal entropy; (c) JS distance between the observed and the bootstrap entropy, for the same datasets.




Figure 3: Individual entropy of subgroup composition for adult spider monkeys in the 2009 dataset (a) and 2013 dataset (b); chimpanzees in the 2008 dataset (c) and 2009 dataset (d); geladas in the 2014 dataset (e) and 2015 dataset (f). Circles and squares represent females and males, respectively. Solid symbols correspond to the observed entropy (H) and empty symbols to the bootstrap entropy (H^b). Labels for each individual in the horizontal axis.